

Division of Laboratory Medicine

Virology

Hepatitis C antibody (HCV) screen and confirmation (Dried Blood Spot)

This service is designed for those patients on whom it is difficult to obtain venous blood, especially for intra venous drug users.

Contact the laboratory for supply of postal packs containing all necessary items for using this service.

Paediatric/infant packs are available on request

General information

Collection container (including preservatives):

Envelope with desiccant

Manchester Medical Microbiology Partnership Grade 226 Blood Collection Card	1	00000
Tuck Cover Here		PerkinElmer 226 LOT 102277 / 314470
	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Tuck Cover Here
Name	· ·	Name
Date of Birth	-	Date of Birth

Specimen type: Dried blood spot card

Collection:



Division of Laboratory Medicine

Virology

Sample Collection Disposable gloves should be worn for the whole procedure to protect you and to protect the sample from degradation. Safely discard the lancet in a sharps container. Ensure the patients hands are clean. If the hands are cold ask the patient to rub them together. It is important the puncture site is warm. 3) Identify and clean the puncture site: - Use one of the outer 3 fingers - Avoid finger pad and nail bed - Clean site with alcohol wipe th 8) Spot blood onto each of the Spot blood onto each of the 5 circles on the card. To en-sure enough blood flow: - Hold the puncture site down - Apply intermittent pressure Avoid strong repetitive pressure (milking), it can damage the area. NOTE: FILL each circle with 4) Break the Seal by Twisting blood. Failure to do so may reduce the sensitivity of the screening tests. • 5) Hold the lancet with two fingers and place at the puncture site. 9) Leave the card to dry for a minimum of 30 minutes 10) Place card along with one bag of desiccant in the plastic carrier attached to the request from. 11) Ensure the required details have been entered on both the request form and the dried blood spot card. Place the re-quest card in the envelope pro-vided and send back to the laboratory. 6) Gently apply pressure until the lancet is activated. A click sound will be heard when this has happened. An

Specimen transport: No special needs

Minimum volume of sample: 7mm diameter blood spot

Special precautions: Ensure blood spot has dried and submit with desiccant pouch enclosed

Laboratory information

Measurement units: Not applicable

Biological reference units: Not applicable

Turnaround time for provisional result (working days): 5 days for screen Turnaround time to Final result (working days): 7 days for confirmation

Clinical information

Clinical decision points: Not applicable

Factors known to significantly affect the results: Spots too small, not all spots filled with blood

(Last updated March 2023)