

Laboratory Medicine Care Division

Immunology

Farmer's lung precipitins

General information

These tests detect the presence of IgG antibodies to *Micropolysporium faeni* (Farmer's lung precipitins). Please note from April 2022 the laboratory no longer tests for *Thermoactinomyces vulgaris*.

Specimen transport: At room temperature

Repeat frequency: At significant change of clinical symptoms

Special precautions: None

Laboratory information

Normal reference range:

- *Micropolysporium faeni* - 0 - 22 mg/L

Volume and sample type: 4ml serum

Method: FEIA (Fluoroenzyme Immunoassay)

Turnaround time (calendar days from sample receipt to authorised result): Median - 4

Participation in EQA Scheme: UK NEQAS for FUNGAL and RELATED ANTIGENS

Clinical information

Indications for the test: Hypersensitivity to *Micropolysporium faeni* is a common cause of Farmer's lung, hypersensitivity pneumonitis occurring 12 to 24 hours after exposure to mouldy hay.

Factors affecting the test: In some cases of farmer's lung, hypersensitivity to other fungal species may be implicated.

The absence of IgG to *Micropolysporium faeni* does not rule the diagnosis out. These antibodies can be found in some healthy individuals. The presence of these antibodies supports, but does not confirm, a diagnosis of farmer's lung. There are numerous other causes of hypersensitivity pneumonitis including other fungal spores and occupational antigens. Please discuss with the laboratory if these may be incriminated.

(Last updated August 2026)