




# Understanding your child's heart

## Tricuspid atresia



## Contents

---

About this booklet	5
The normal heart	6
What is congenital heart disease?	8
What causes congenital heart disease?	8
Why me? Why my child?	9
What is tricuspid atresia?	10
What are the symptoms of tricuspid atresia?	12
What other conditions are associated with tricuspid atresia?	12
How is tricuspid atresia diagnosed?	13
How is tricuspid atresia treated?	14
What happens as my child grows up?	19
The specialist centre for congenital heart disease	20
Coping with everyday life	22
General advice for the future	25
What is the risk of having another child with congenital heart disease?	27
Support	28
About the British Heart Foundation	29
References	31
Contacts	32
Hospital visits	33
The medical terms and what they mean	34
Index	36
Have your say	39



## About this booklet

---

This booklet is written for the parents of babies and children with tricuspid atresia, and for their relatives and friends. It explains:

- what tricuspid atresia is and how it is diagnosed
- how tricuspid atresia is treated
- the benefits and risks of treatments
- how to cope as a parent of a child with tricuspid atresia
- where to go for more support.


This booklet does not replace the advice that doctors or nurses may give you, but it should help you to understand what they tell you.

## The normal heart

The heart is a muscular pump which pumps blood through the body and lungs. There are four chambers in the heart. The two upper ones are called the right atrium and left atrium. These are separated by a wall called the atrial septum. The two lower chambers are called the right and left ventricles, and are separated by a wall called the ventricular septum. See the illustration opposite.

On each side of the heart, blood passes from the atrium, through a heart valve – the tricuspid valve on the right, and the mitral valve on the left – into the ventricle. The ventricles are the main pumping chambers of the heart. Each ventricle pumps blood out into an artery. The right ventricle pumps blood – blue in the illustration – into the pulmonary artery (the blood vessel that takes blood to the lungs). The left ventricle pumps blood – red in the illustration – into the aorta (the blood vessel that takes blood to the rest of the body). Blood flows from the right side of the heart, through the pulmonary valve into the pulmonary artery, and then to the lungs where it picks up oxygen. The oxygen-rich blood flows back into the left side of the heart through the pulmonary veins. The left ventricle then pumps the oxygen-rich blood out of the heart through the aortic valve and into the aorta, and all around the body. The blood then returns to the right side of the heart through two main veins – one from the upper body (superior vena cava), and the other from the lower body (inferior vena cava).

## The normal heart



## What is congenital heart disease?

---

Congenital heart disease is an abnormality of the heart that developed in the womb. In some cases, congenital heart disease is diagnosed when the baby is still developing in the womb, but in most cases the problem is not discovered until after the baby is born. There are lots of different types of congenital heart disease. *Tricuspid atresia* is a very serious type.

## What causes congenital heart disease?

---

We know that, in most cases of congenital heart disease, something has gone wrong in the early development of the fetus, at the very early stages of the pregnancy. In most cases, we don't understand why the baby's heart did not develop normally. In some cases, congenital heart disease can be part of a syndrome that the baby is born with. (A syndrome means a group of symptoms that appear together.)

## Why me? Why my child?


---

It is not unusual for parents of children with congenital heart disease to blame themselves or to be angry. Anger, disappointment, fear and guilt are all normal feelings to have when you are told that there is something wrong with your child. At first it may be difficult to cope with and it can take a while for the news to sink in.

Many pregnant women or mothers, and their partners, ask themselves what they did during their pregnancy that could have caused their baby's heart to develop with heart disease. But the reality is that it can happen to anyone. In fact congenital heart disease happens in about 7 in every 1,000 pregnancies.<sup>1</sup> For more than half of these children, the heart disease is only a minor problem which either doesn't need any treatment, or which can be successfully corrected with surgery. problem which either doesn't need any treatment, or which can be successfully corrected with surgery. For others it is more serious and, sadly, some children don't survive. However, thanks to advances in diagnosis and treatment, most children can be helped to have a good quality of life.

## What is tricuspid atresia?


### The normal heart



Tricuspid atresia is a very serious type of congenital heart condition. There are three main abnormalities:

- The tricuspid valve failed to develop, so there is no connection between the right atrium and the right ventricle. Instead, blood flows to the left atrium through a hole in the atrial septum.

### Tricuspid atresia



- The right ventricle is very small.
- There is a hole in the ventricular septum – the wall between the two ventricles. This is called a '**ventricular septal defect**'.

In the normal heart, blood flows from the right atrium through the tricuspid valve to the right ventricle, and from there it goes through the pulmonary artery to the lungs. (See the illustration on page 10.) In children with tricuspid atresia, the blood cannot flow from the right atrium into the right ventricle. Instead, it flows from the right atrium to the left atrium through a hole in the atrial septum. From the left atrium, the blood flows to the left ventricle, which pumps blood to both the aorta and (through the ventricular septal defect) to the pulmonary artery. (See illustration on page 11).

### **What are the symptoms of tricuspid atresia?**

---

The symptoms of tricuspid atresia vary from one child to another, depending on exactly which abnormalities they may have. Some children look remarkably well, some children become blue if there is too little blood flowing to the lungs, while others become breathless early in life if there is too much blood flowing to the lungs. Children who are breathless may not gain weight normally.

### **What other conditions are associated with tricuspid atresia?**

---

Children with tricuspid atresia may also have other abnormalities of the heart. In some children, the main arteries are 'transposed'. This means that the aorta arises from the right ventricle instead of the left, and the

pulmonary artery arises from the left ventricle instead of the right. In some children there may be a narrowing in the aorta, known as **coarctation of the aorta**. Others may have **pulmonary stenosis** (narrowing of the pulmonary valve), or **pulmonary atresia** (complete blockage of the pulmonary valve). So, some children will have too much blood flowing to the lungs, and some will have too little. Your cardiologist will discuss your child's individual condition with you.

### **How is tricuspid atresia diagnosed?**

---

In some cases, tricuspid atresia may be detected before birth, but in other cases it is not diagnosed until after the baby is born. Usually, the only test that is needed to make the diagnosis is an **echocardiogram**. This is an ultrasound scan of the heart. It is very similar to the scans that are carried out during pregnancy. It doesn't hurt your baby at all.

## How is tricuspid atresia treated?

It is not possible to correct tricuspid atresia with surgery, but there are operations which can help children to have a reasonable quality of life.

Surgery is carried out in stages as your child grows. The type and timing of surgery recommended for an individual child with tricuspid atresia will depend on which abnormalities they have. Some children don't need any surgery until they are a few years old, but most will need some treatment as young children.


### Initial surgery

Children who have too much blood flowing to the lungs need surgery to reduce the blood flow. This is called **pulmonary artery banding**. Children with too little blood flowing to the lungs need surgery to increase the flow. This is called a **shunt operation**. We explain more about these treatments below. Children with coarctation of the aorta will also need to have this repaired. Your cardiologist will explain what type of treatment is recommended for your child.

### Pulmonary artery banding

This is an operation to reduce the blood supply to the lungs. The timing of the operation depends on how breathless the baby becomes. Sometimes it is needed within the first few months of life.

## Tricuspid atresia with a pulmonary artery band





This surgery involves placing a band around the pulmonary artery to narrow it. This is done without using a heart-lung bypass machine, but your baby will be given a general anaesthetic. The surgery usually leaves a scar at the side of the chest rather than in the middle. The band reduces the high blood flow to the lungs, reducing breathlessness and lowering the blood pressure in the pulmonary artery, to try to prevent lung damage. (See the illustration on page 15).


Your baby will need to stay in hospital for several days after the operation. You will need to take your baby to the outpatients department a few weeks after the operation, to see the paediatric cardiologist for a check-up.

### Arterial shunt operation

For this operation your child will need to have a general anaesthetic. The operation does not usually involve the use of a heart-lung machine. An arterial shunt operation involves placing a small tube made of synthetic material between a branch of the aorta and the pulmonary artery. (See the illustration opposite).

The surgery usually leaves a scar at the side of the chest rather than in the middle. The shunt allows more blood to flow to the lungs and makes the baby pinker.

### Tricuspid atresia with a shunt



Your baby will need to stay in hospital for several days after the operation. You will need to take him or her to the outpatients department a few weeks after the operation, to see the paediatric cardiologist for a check-up.

### **What risks are involved with pulmonary artery banding or an arterial shunt operation?**

About 96 of every 100 children who have pulmonary artery banding or an arterial shunt operation survive the surgery.<sup>2</sup> However, a small proportion of these will need another operation quite soon after the first one, either to tighten the band or to enlarge the shunt.

### **Further surgery**

The exact nature and timing of any further surgery will depend on your child's progress. The type of surgery most commonly carried out is a **cavopulmonary shunt**, which involves connecting the superior vena cava directly to the pulmonary arteries. Further surgery after this usually involves redirecting the blood flow from the inferior cava to the pulmonary artery. This is called a **total cavopulmonary connection**, or **TCPC** for short. For more information on these treatments, see our booklet *Understanding your child's heart: Single ventricle circulation*.

Your cardiologist will explain what type of treatment is recommended for your child.

## **What happens as my child grows up?**

---

Tricuspid atresia is a complex condition, and even with surgical treatment it cannot be corrected. Although surgery can give a reasonable quality of life, the operations are relatively new so we do not know how long children with this condition will live for. The longest survivors at present are in their 30s. Heart transplantation may be an option for some patients, although this is rarely considered before adulthood.

Children with tricuspid atresia are almost always limited to some extent in their physical activities, but specific restrictions on exercise are usually not necessary. It is usually best to allow children with this condition to join in with their friends' activities, including some sport at school, and for them to judge for themselves what they are able to do.

Whatever type of treatment your child has for tricuspid atresia, regular check-ups with a cardiologist are very important – not just in childhood, but throughout adult life too.

## The specialist centre for congenital heart disease

---

Your child will continue to have check-ups at a specialist centre for congenital heart disease throughout their life. Up to the age of about 16, he or she will go to a specialist centre for children with congenital heart disease (paediatric cardiac centre).

If the specialist centre is quite a distance from your home, it may be possible to make arrangements for your child to be looked after under a 'shared care' system (some general hospitals have paediatricians who have had special training in cardiology). This means that your child will be looked after locally, but will go to the specialist centre for specialised treatment. (See page 24 for information about claiming travel expenses for visits to the specialist centre.)

### The specialist team

At the specialist centre, a large multidisciplinary team of people will be involved in caring for your child and your family's needs. (Multidisciplinary means that it includes several different types of health professionals.) The team usually consists of:

- a consultant paediatric cardiologist
- a specialist registrar (a doctor who is specifically training in children's heart problems)
- a cardiac nurse specialist or cardiac liaison nurse
- a consultant paediatric surgeon or paediatric cardiothoracic surgeon
- cardiac technicians

- a paediatric physiotherapist
- a paediatric dietitian
- ward-based paediatric nurses
- a social worker
- an occupational therapist
- a speech and language therapist, and
- a play specialist.

All of these people are involved in planning the care for each patient. You probably won't need to see all of them, but it is important to know who is there and available to help you with any problems you may come across. For example, the social worker can be a very useful source of information on what benefits you might be entitled to claim, and whether you can claim your travel expenses for visiting your child in hospital, or for visits to the specialist centre. (For more on this, see page 24.)

The specialist centre will also have access to psychology services which you can use to help your child or your family if you need help at difficult times.

### Specialist adult congenital heart disease centres

When your child grows into adulthood, it is important that he or she carries on going to a specialist centre for check-ups. There are several specialist centres in the UK for adults with congenital heart disease. Your child's care will be transferred to an adult specialist centre usually at around the age of 16. These centres usually have a multidisciplinary team with the same mix of professionals as in the children's specialist centre

(see page 20). When your child is nearing adulthood, your specialist paediatric centre will start preparing you and your child for the move to the adult specialist centre, to make the transition as smooth as possible.

## **Coping with everyday life**

---

Having a child with a heart condition in hospital can be very difficult for a number of reasons. You are having to care for your child in hospital, and may be anxious about the treatment your child has to have or how well he or she is recovering. You might also be worrying about being away from home, or about your other children and who's looking after them, or about your work or your finances.

Hospital staff recognise the stress that you and your family might be under. Ask the nurse who is looking after your child about the support services available within the hospital – such as psychology services, welfare rights advisers and social workers. Below we give a brief guide to the benefits and help you may be able to get.

### **Financial issues**

Financial problems may arise because you need to stay in hospital with your child. The hospital may be a long way from home, and you may not know how long you will need to stay there for. This can affect your finances, as you may have to take time off work and will have extra costs such as travel expenses and buying meals while in hospital. This can be an extra anxiety, and difficult to cope with.

If you are worried about your finances, it is important to discuss your situation with a hospital social worker or cardiac liaison nurse, or with the Citizens Advice Bureau. They may be able to advise you on the benefits you can claim. Also, an organisation called Working Families can give you advice on the phone – on 0800 013 0313 – about your rights as an employee if you need to take time off work to be with your child.

### **Low-income benefits**

Benefits for people on a low income include Income Support, housing and council tax benefits, and Tax Credits. If your income goes below a certain amount, you may be able to claim benefits. However, you have to meet certain criteria in order to get these benefits. The criteria vary from one benefit to another, so you should get specialist advice from a hospital social worker, Citizens Advice Bureau or Jobcentre Plus.

### **Disability benefits**

Some children with congenital heart disease will qualify for a *Disability Living Allowance* (DLA), but most will not. Ask the specialist nurse or social worker for advice. If your child needs extra care because of his or her condition, you may be able to apply for this benefit. There is a mobility and a care component to the benefit. It can be difficult to get Disability Living Allowance for a child. You will need to show that your child needs more attention or supervision than other children of the same age. To get a claim form for Disability Living Allowance, call 0800 88 22 00. Or you can get a form from your GP or your local Jobcentre Plus office, or claim online at [www.direct.gov.uk](http://www.direct.gov.uk)

## Carer's Allowance

You may be entitled to *Carer's Allowance* if your child is awarded the middle or higher care component of Disability Living Allowance. However, you will only get Carer's Allowance if your earnings are below a certain limit. To get a claim form for the Carer's Allowance, call 01253 856 123. Or you can get a claim form from your local Jobcentre Plus, or claim online at [www.direct.gov.uk](http://www.direct.gov.uk)

## Fares to hospital

If you get Income Support, or have a Tax Credit exemption card, or are assessed as being on a low income, you are entitled to get reimbursed for your fares to hospital appointments and inpatient treatment. If you think you might qualify because of your income, complete a form HC1. To get one, call 0845 850 1166. The forms are usually also available from Jobcentre Plus offices and NHS hospitals.

## Fares for visiting your child in hospital

If you are getting Income Support, income-based Jobseeker's Allowance, or Pension Credit, you may be able to get help towards the cost of fares for hospital visiting, from the Social Fund. You should apply for a Community Care Grant on form SF300, which is available from your local Jobcentre Plus office or from the website of the Department for Work and Pensions at [www.dwp.gov.uk](http://www.dwp.gov.uk). However, these payments are not given in all cases.

**The benefits system is very complex, so it is important to get specialist advice on what you may be entitled to, from the hospital social worker or Citizens Advice Bureau.**

## General advice for the future

---

### Endocarditis

Everyone who has tricuspid atresia is at risk of getting *infective endocarditis*, both before and after surgery or treatment. Infective endocarditis is a rare condition where the inner lining of the heart, most commonly one of the heart valves, becomes infected.

Infective endocarditis is a serious condition which can be life-threatening if it's not treated quickly. Nowadays, if it is diagnosed early, most people with it will recover well with antibiotic treatment, although some damage may occur to the heart valves as a result of the infection.

Endocarditis is caused by a bacterium, or (rarely) another type of infective organism that is in the bloodstream, which settles onto the abnormal structure or defect in the heart. Although it is not possible to prevent all bacteria from getting into the bloodstream, there are some things your child can do, as he or she grows up, to reduce the risk of getting endocarditis:

- Maintain good oral hygiene and have regular check-ups with a dentist
- Avoid body piercing and tattooing
- Never inject recreational drugs.

If your child develops flu-like symptoms with a temperature **which persist for over a week**, you should visit your GP as your child may need a blood test. Make sure that the GP knows that

your child is at increased risk of getting endocarditis. You can do this by showing the GP an *Endocarditis warning card*. You can get this card from the British Heart Foundation (BHF) by calling either the Heart HelpLine on **0300 330 3311** (local rate number) or the BHF Orderline on **0870 600 6566**.

### **Pregnancy**

If you have a daughter with congenital heart disease, you need to be aware that pregnancy could carry risks to both the mother and the baby. So, when your daughter gets older, it's particularly important that she avoids having an unplanned pregnancy. You will need to discuss this with your daughter in whichever way you, as a parent, think is appropriate for her. If your daughter wants to have a baby, it's best that she speaks to her cardiologist about it first, so that the pregnancy can be planned for when your daughter's heart condition is most stable.

People who have congenital heart disease themselves have an increased risk of having a child with a heart problem. This applies to both males and females. You can discuss this with your cardiologist. Early scans in pregnancy can be arranged to look for heart disease in the baby.

## **What is the risk of having another child with congenital heart disease?**

---

If you have one child with congenital heart disease, there is about a 1 in 50 chance of having another child with congenital heart disease.<sup>3</sup> However, this risk may be higher (or lower) depending on the type of congenital heart disease your child has. Because your risk of having another child with congenital heart disease is higher than it is for other people, you may be offered a special scan at an early stage in future pregnancies, to look at the baby's heart. Ask your midwife or GP for more information on having a scan earlier than usual.

If you have had two children with congenital heart disease, the risk of having another child with heart disease rises to about a 1 in 10 chance. This may sound like a high risk, but you still have a much better chance of the baby's heart being normal than abnormal. If there is a recurrence, the heart disease may not always be of the same type.<sup>3</sup>

## Support

---

The following support groups may be able to offer you further information, advice and support:

### **Little hearts matter**

11 Greenfield Crescent, Edgbaston, Birmingham,  
West Midlands, B15 3AU  
Phone: 0121 455 8982. Website: [www.lhm.org.uk](http://www.lhm.org.uk)

### **Action for Sick Children**

32b Buxton Road, High Lane, Stockport SK6 8BH  
Phone: 0800 074 4519. Website: [www.actionforsickchildren.org](http://www.actionforsickchildren.org)

### **Children's Heart Federation**

Level One, 2-4 Great Eastern Street, London EC2A 3NW  
Phone: 0808 808 5000. Website: [www.childrens-heart-fed.org.uk](http://www.childrens-heart-fed.org.uk)

### **Grown Up Congenital Heart Patients Association**

Saracen's House, 25 St Margaret's Green, Ipswich IP4 2BN  
Phone: 0800 854759. Website: [www.guch.org.uk](http://www.guch.org.uk)

### **www.yheart.net**

A website for young people with heart conditions

## About the British Heart Foundation

---

The British Heart Foundation is the nation's heart charity, saving lives through pioneering research, patient care and vital information.

### **What you can do for us**

We rely on donations to continue our vital work. If you would like to make a donation to the BHF, please ring our Supporter Services team on **0844 847 2787** or contact us through our website at [bhf.org.uk/donate](http://bhf.org.uk/donate) or send it to us at the address on the back cover.

## For more information

---

### **British Heart Foundation website**

[bhf.org.uk](http://bhf.org.uk)

For up-to-date information on heart disease, the BHF and its services.

### **Heart HelpLine**

**0300 330 3311** (local rate number)

For information and support on anything heart-related.

## Other resources

---

### Understanding your child's heart series

This booklet is one of the booklets in the *Understanding your child's heart* series. For a full list of the booklets available in this series, see our website [bhf.org.uk/congenital](http://bhf.org.uk/congenital) or call the Heart Helpline on **0300 330 3311** (local rate number).

### Children with congenital heart disease (DVD)

Three families share their experiences from diagnosis to treatment, and staff at the Evelina Children's Hospital offer guidance on parents' common concerns.

### Operation Fix-it

A short story book about eight-year-old Tom's experience in hospital for a heart operation. Prepares children for their hospital visit in an interesting and sometimes humorous way.

### To order any of our resources

- call the BHF Orderline on **0870 600 6566**
- email [orderline@bhf.org.uk](mailto:orderline@bhf.org.uk) or
- visit [bhf.org.uk/publications](http://bhf.org.uk/publications)

You can also download many of our publications from our website. For information on other BHF booklets and DVDs ask for a copy of the *Our heart health catalogue*.

## References

---

1. Petersen S, Peto V, Rayner M. 2003. *Congenital Heart Disease Statistics 2003*. Oxford: British Heart Foundation Health Promotion Research Group.
2. Specific procedures national data. The congenital heart disease website. London: The Information Centre. Accessed from: [www.ccad.org.uk](http://www.ccad.org.uk)
3. Gill HR, Splitt M, Sharland GK, Simpson JM. 2003. Patterns of recurrence of congenital heart disease: An analysis of 6,640 consecutive pregnancies evaluated by detailed fetal echocardiography. *Journal of the American College of Cardiology*; 42: 923-9.





## The medical terms and what they mean

---

<b>aorta</b>	The main artery of the heart. It supplies oxygen-rich blood to the body.
<b>atrium</b>	One of the two upper chambers of the heart.
<b>cardiac</b>	To do with the heart.
<b>cardiologist</b>	A consultant specialising in heart disease.
<b>coarctation of the aorta</b>	Narrowing of the aorta.
<b>congenital</b>	From birth.
<b>echocardiogram</b>	An ultrasound scan used to produce pictures of the heart and blood vessels.
<b>endocarditis</b>	Infection of the lining of the heart or its valves.
<b>genetic</b>	To do with the information that is passed from parents to children through genes in sperm and eggs.
<b>heart-lung machine</b>	A machine that pumps blood around the body while the heart is stopped during an operation.

<b>paediatric</b>	To do with paediatrics – the study of children's diseases.
<b>pulmonary</b>	To do with the lungs.
<b>pulmonary stenosis</b>	When the pulmonary valve is very narrow.
<b>septum</b>	The wall that keeps the right and left sides of the heart separate.
<b>stenosis</b>	See pulmonary stenosis.
<b>ventricle</b>	One of the two lower chambers of the heart.
<b>ventricular</b>	To do with the ventricle or ventricles. (See above.)
<b>ventricular septal defect</b>	A hole between the two ventricles of the heart. Also called VSD.

## Index

---

aorta	6, 7
aortic valve	6, 7
arterial shunt operation	15
atrium	6, 7
benefits	23
blueness	12
Carer's Allowance	24
causes of congenital heart disease	8
cavopulmonary shunt	18
centre: specialist centre	20, 21
coarctation of the aorta	13
congenital heart disease	8
diagnosis	13
Disability Living Allowance	23
echocardiogram	13
endocarditis	25
fares to hospital	24
finance	22
future	19, 25
heart: how the normal heart works	6
infection	25
long term outlook	19
mitral valve	6, 7
money	24
normal heart	6
operation	14–18

pregnancy	26
professionals	20, 21
pulmonary artery	6, 7
pulmonary artery banding	14
pulmonary stenosis	13
pulmonary valve	6, 7
pulmonary veins	6, 7
repair of coarctation of the aorta	14
risks of surgery	18
septum	6, 7
specialist centre	20, 21
specialist team	20, 21
stenosis	13
support groups	28
surgery	14–18
symptoms	12
TCPC	18
team	20, 21
total cavo-pulmonary connection	18
travel expenses	24
treatment	14
tricuspid valve	6, 7
valves	6, 7
ventricle	6, 7
ventricular septal defect	11



## Have your say

We would welcome your comments to help us produce the best information for you. Why not let us know what you think? Contact us through our website at [bhf.org.uk/contact](https://www.bhf.org.uk/contact) Or, write to us at the address on the back cover.

## Acknowledgements

The British Heart Foundation would like to thank:

- Ms Carole Wingett, Social Worker, Royal Brompton Hospital.

Particular thanks for their work on this series of booklets are due to:

- Dr Gurleen Sharland, Consultant Fetal Cardiologist, Evelina Children's Hospital, Guy's & St Thomas' NHS Foundation Trust
- Dr James Gnanapragasam, Consultant Paediatric Cardiologist, Southampton General Hospital
- Dr John Gibbs, Consultant Paediatric Cardiologist, Leeds General Infirmary.

# HEART HELPLINE

For information and support on anything heart-related



**0300 330 3311**

local rate number



**bhf.org.uk**

Phone lines open 9am to 6pm Monday to Friday



## British Heart Foundation

Greater London House  
180 Hampstead Road  
London NW1 7AW  
Phone: 020 7554 0000  
Fax: 020 7554 0100  
Website: [bhf.org.uk](http://bhf.org.uk)

C10/0210