



Pain Relief in Labour



This information will give you some idea about the pain of labour and giving birth, and what can be done to make it less painful. The people who are looking after you (for example, your midwife, anaesthetist or obstetrician) will give you more information about the types of pain relief that your own hospital or birth centre provides. We hope that, if you know what to expect and what pain relief is available, giving birth to your baby will be a satisfying

experience.

Throughout this information, we have used references to show where we have got our information. We have listed these at the end.

What will labour feel like?

- While you are pregnant, you may feel your uterus (womb) tightening from time to time. These are called Braxton Hicks contractions. When you go in to labour, this tightening feeling becomes regular and much stronger.
- The tightening may cause pain that feels like period pain, and usually becomes more painful the further you get in to labour. Different women experience labour pains in different ways.
- Usually, your first labour will be the longest.
- If medication is used to start off (induce) labour or speed up your labour, your contractions may be more painful.
- Most women use a range of ways to cope with labour pain (see reference 1). It is a good idea to have an open mind and be flexible.

Preparing for Labour

Antenatal parenting classes help you prepare for the birth. These classes are run by midwives and by other organisations that support people in being parents and giving birth. The classes will help you understand what will happen in labour and may help you to feel less anxious.

At antenatal classes, the midwife will tell you what is available to reduce labour pain. If you need more information about epidurals (an injection into your back to numb the lower half of your body), the midwife can arrange for you to meet an anaesthetist to talk about this. If you cannot go to antenatal classes, you should still ask your midwife about what is available to reduce the pain. You can then discuss this with the midwife who cares for you while you are in labour.

Where you choose to give birth can affect how painful it is. If you feel at ease in the place you give birth, you may be more relaxed and less anxious about labour (see reference 2). For some women this means giving birth at home, but other women feel reassured by the support offered at a hospital or birth centre. Many hospitals try to make the labour rooms look homely and encourage you to play music you like to help you feel more relaxed.

If you are planning to give birth in a hospital or birth centre, it may be helpful to look round to find out what facilities they have.

Having a friend or birth partner with you while you are in labour can be helpful for you (see reference 3). It is important to talk to your birth partner about your concerns and what you want, and they can help you to focus during the birth.

What pain relief is available?

It is difficult to know beforehand what sort of pain relief will be best for you. The midwife who is with you in labour should be the best person to give you advice. Here is some information about the main methods of pain relief available.

Self help methods

- Breathing calmly may increase the amount of oxygen that is supplied to your muscles, and so make the pain less intense. Also, because you are focusing on your breathing, you are likely to be less distracted by the pain.
- It can be difficult to relax when you are in pain, so it can be helpful to practice before you actually go in to labour. There are a number of different ways you can learn to relax.
- You may find that having a massage while you are in labour can be very comforting and reassuring.



Using a birthing pool during labour

There are not many studies that have looked at the benefits and risks of using a birthing pool. However, it has been shown that if you have your labour in water you will find it less painful and you will be less likely to need an epidural to reduce the pain (see reference 4). There are some concerns that if the water is too warm your baby may show signs of distress during labour, but studies have shown that there is no more risk to you or your baby if you have your labour in water than if you have it out of water. The midwife will continue to monitor your progress and your baby's well being.

Many maternity units have birthing pools, but these may not be available when you need them. It is worth checking with your midwife if there is a pool and whether you would be able to use it.

Complementary therapies (these do not use medications)

Complementary therapies (for example, aromatherapy) may help some women to cope with pain during labour. If you are thinking about using these, it is important that you get advice from a person trained in that therapy. This booklet does not cover homoeopathy (using very dilute ingredients to reduce pain) and herbal remedies (produced from plants).

Aromatherapy involves using concentrated essential oils to reduce fear, improve your well being and give you encouragement.

Reflexology is based on the idea that certain points on your hands and feet relate to points on the rest of your body. We do not know how it works, but it may work in a similar way to acupuncture (see below). A reflexologist usually massages points on your feet that relate to the parts of your body that are painful in labour.

Hypnosis and acupuncture are becoming more popular for some women to help them through labour. Very few maternity units provide these services on the NHS, so you would need to find a qualified therapist before you go into labour.

Hypnosis can distract you from the pain. You can be trained to do the hypnosis yourself (self-hypnosis), and you will need to practise this while you're pregnant. Otherwise, a hypnotherapist will have to be with you while you are in labour.

Acupuncture involves putting needles into points on your body to help reduce the pain. The therapist would need to be with you during your labour. Some studies suggest that women who use these therapies feel in control of their labour and use less medication to reduce pain (see reference 5). However, not all parts of the country have therapists with the necessary level of skill, and their support can be quite expensive.



Transcutaneous Electrical nerve stimulation (TENS)

- A gentle electrical current is passed through four flat pads stuck to your back. This creates a tingling feeling. You can control the strength of the current yourself.
- It is sometimes helpful at the beginning of labour, particularly for backache. If you hire a TENS machine, you can start to use it at home. Some hospitals will also lend you one.
- TENS machines have no known harmful effects on your baby

While you may manage your labour with only the help of TENS, it is more likely that you will need some other sort of pain relief later on in labour.

Entonox

- Entonox is a gas made up of 50% nitrous oxide and 50% oxygen. It is sometimes known as gas and air.
- You breathe Entonox through a mask or mouthpiece
- It is simple and quick to act, and wears off in minutes.
- It sometimes makes you feel light-headed or a little sick for a short time
- It does not harm your baby and it gives you extra oxygen, which may be good for you and your baby.
- It will not take the pain away completely, but it may help.
- You can use it at any time during labour.

You control the amount of Entonox you use, **but to get the best effect it is important to get the timing right**. You should start breathing Entonox as soon as you feel a contraction coming on, so you will get the full effect when the pain is at its worst. You should not use it between contractions or for long periods as this can make you feel dizzy and tingly. In some hospitals, other substances may be added to Entonox to make it more effective, but these may make you sleepier.

Opioids



Opioids are painkillers which work in a similar way to morphine. They include painkillers such as pethidine, as well as diamorphine (which is being used more and more in the United Kingdom). Research has shown that women who receive IM diamorphine are more likely to be satisfied with their analgesia (pain relief) when compared with IM Pethidine (see reference 6). ('IM' stand for 'intramuscular' and means that the drug is injected into the muscle.) Examples of other opioids include morphine meptizol, fentanyl and remifentanyl. All these morphine-like painkillers act in a similar way.

- A midwife usually gives opioids by injecting them into a large muscle in your arm or leg
- The pain relief is often limited. You will start to feel the effects after about half an hour and they may last a few hours.
- Opioids are less effective at easing pain in labour than Entonox.
- Although pain relief may be limited, some women say it makes them feel more relaxed and less worried about the pain (see reference 7)
- Other women are disappointed with the effect of opioids on their pain and say they feel less in control.

Side effects of opioids

- They may make you feel sleepy.
- They may make you feel sick, but you will usually be given anti-sickness medication to stop this
- They delay your stomach emptying, which might be a problem if you need a general anaesthetic
- They may slow down your breathing. If this happens, you may be given oxygen through a face mask and have your oxygen levels monitored.
- They may make your baby slow to take their first breath, but your baby can be given an injection to help with this
- They may make your baby drowsy, and this may mean that they cannot feed as well as normal (especially if you are given pethidine).
- If you are given opioids just before you give birth to the baby, the effect on your baby is very small.

Patient controlled analgesia (PCA)

Opioid can also be given direct into a vein for a faster effect, using a pump that you control yourself by pressing a button attached to the pump. PCA is available in some hospitals if an epidural (an injection into your back to numb the lower half of your body) is not possible or you do not want one.

PCA allows you to give yourself small doses of opioids when you feel that you need them. You have control over the amount of opioid you use. For safety reasons, the PCA limits how quickly you take the opioid. However, if you use the PCA for a long time, some opioids may build up in your body which may increase the side effects of the opioid on you and your baby.

In a few maternity units, you may be offered PCA using an opioid called remifentanyl (see references 8 and 9). Your body breaks down remifentanyl very quickly, so the effects of each dose do not last long. This opioid has a strong effect on pain but it is also more likely to slow down your breathing, so your breathing needs to be checked carefully. However, its effects can be reversed quickly and it will not affect your baby.

Epidurals and spinals

- Epidurals and spinals are the most complicated method of pain relief and are carried out by an anaesthetist.
- An anaesthetist is a doctor who is specially trained to provide pain relief and drugs that make you go to sleep. Pain relief during operations can be provided using general anaesthesia, epidurals or spinals. For more information on these types of anaesthesia for a Caesarean section, please take a look at [Caesarean section information](#) on LabourPains.com
- Epidurals and spinals are the most effective method of pain relief.
- For an epidural, the anaesthetist inserts a needle into the lower part of your back and uses it to place an epidural catheter (a very thin

tube) near the nerves in your spine. The epidural catheter is left in place when the needle is taken out so you can be given painkillers during your labour. The painkillers may be a local anaesthetic to numb your nerves, small doses of opioids, or a mixture of both.

- An epidural may take 40 minutes to give pain relief (including the time it takes to put in the epidural catheter and for the painkillers to start working).
- An epidural should not make you feel drowsy or sick.
- Having an epidural increases the chance that your obstetrician will need to use a ventouse (a suction cap on your baby's head) or forceps to deliver your baby.
- An epidural can usually be topped up to provide pain relief if you need a ventouse, forceps or a Caesarean section.
- An epidural will have hardly any effect on your baby.

Spinal and combined spinal-epidural (CSE)

Epidurals are rather slow to act, especially if you have one late in labour. If the painkillers are given direct into the bag of fluid surrounding the nerves in your back, they work much faster. This is called a spinal. Unlike an epidural, it is given as a one-off injection without a catheter. If an epidural catheter is put in at the same time, this is called a combined spinal-epidural.

In some hospitals, a combined spinal-epidural is given to almost all women who want strong pain relief instead of an epidural. In others, a combined spinal is only used for a small number of women.

Who can and cannot have an epidural?

Most people can have an epidural, but certain medical problems (such as spina bifida, a previous operation on your back or problems with blood clotting) may mean that it is not suitable for you. The best time to find out about this is before you are in labour. If you have a complicated or long labour, your midwife or obstetrician may suggest that you have an epidural as it may help you or your baby. If you are overweight, an epidural may be more difficult and take longer to put in place. However, once it is in you will have all the benefits.

What does an epidural involve?

First, a cannula (a fine plastic tube) will be put in a vein in your hand or arm, and you will usually have a drip (intravenous fluid) running as well (you may also need a drip in labour for other reasons, such as to give you medication to speed up your labour or if you are being sick). Your midwife will ask you to curl up on your side or sit bending forwards, and your anaesthetist will clean your back with an antiseptic.

Your anaesthetist will inject local anaesthetic into your skin, so that putting in the epidural does not usually hurt much. The epidural catheter is put into your back near your nerves in the spine. Your anaesthetist has to be careful to avoid puncturing the bag of fluid that surrounds your spinal cord, as this may give you a headache afterwards. It is important to keep still while the anaesthetist is putting in the epidural, but after the epidural catheter is fixed in place with tape you will be free to move.

Once the epidural catheter is in place, you will be given painkillers through it. It usually takes about 20 minutes to set up the epidural and 20 minutes for it to give pain relief. While the epidural is starting to work, your midwife will take your blood pressure regularly. Your anaesthetist will usually check that the epidural painkillers are working on the right nerves by putting an ice cube or cold spray on your tummy and legs and asking you how cold it feels. Sometimes, the epidural doesn't work well at first and your anaesthetist needs to adjust it, or even take the epidural catheter out and put it in again.

During labour, you can have extra doses of painkillers through the epidural catheter either as a quick injection (a top-up), a slow, steady flow using a pump, or with a patient controlled epidural analgesia (PCEA) pump, with patient controlled epidural analgesia, you can give yourself doses of the painkiller when you need them by pressing a button attached to the pump. In each hospital there will usually only be one, or possibly two of these methods for keeping the epidural pain relief going.

After each epidural top-up, the midwife will take your blood pressure regularly in the same way as when the epidural was started.

The aim of the epidural is to take away the pain of contractions. Usually, the epidural also completely takes away the pain when your baby is delivered. Some women prefer to have some feeling during the delivery so they have a better idea of how to push the baby out. The epidural cannot be adjusted exactly, so if you want to have some feeling when your baby is delivered, there is more chance that you may have an uncomfortable sensation during labour as well.

Nowadays it is usually possible to reduce the pain of labour without making the lower part of your body very numb or making your legs feel weak. This modern method is called a 'mobile epidural'. You will be able to breastfeed your baby after the epidural.

What if I need an operation?

If you need a Caesarean section, the epidural is often used instead of a general anaesthetic. A strong local anaesthetic is injected into your epidural catheter to make the lower half of your body very numb for the operation. This is safer than a general anaesthetic for you and your



baby. Occasionally the epidural may not work well enough to be used for a Caesarean section. This can happen in 1 in 8 to 10 people. If this happens, you may also need another anaesthetic such as a spinal or general anaesthetic.

If you need a Caesarean section but you do not already have an epidural, a spinal will often be used but with a bigger dose of local anaesthetic than the dose which is used for a spinal in labour.

For more information on epidurals and spinals for a Caesarean section, please take a look at [Caesarean section information](#) on LabourPains.com

Benefits and risks of epidurals

How do we get our facts?

We get our facts from randomised studies and from observational studies.

Randomised studies are when women have either one type of treatment or another, and the effects of the different treatments are compared. Which of the two treatments each woman has is decided randomly (that is, like tossing a coin). The studies usually compare women who have an epidural with women who used other painkillers (such as opioids or Entonox) during labour.

Reference 10 is a review of all the published randomised studies on epidurals in labour. It was carried out by the Cochrane database, which is an independent scientific organisation. The effects of epidurals that we talk about below are from this review, unless we give a different reference. In a few randomised studies, all the women have had an epidural, but the amount of opioid that is used in the epidural is decided randomly.

Observational studies look at large numbers of women who have had an epidural to see what happens during the epidural and afterwards. This is the only way to find out the risk of very rare events.

The following information is based on the results of randomised studies.

Benefits of having an epidural

- Epidurals reduce the pain of labour more than any other treatment.
- With an epidural, there is less acid in the newborn baby's blood (see reference 11).
- With an epidural, there is less need to use medication to make your baby start breathing when they are born, compared with opioids given in other ways (into a muscle or a vein).

Things an epidural does not make a difference to

- With an epidural, you do not have a higher chance of needing a Caesarean section.
- There is no greater chance of long-term backache. Backache is common during pregnancy and often continues afterwards. You may have a tender spot in your back after an epidural which, rarely, may last for months (see reference 12).

Risks while the epidural is being used

- With an epidural, the chance of the obstetrician having to use a ventouse or forceps to deliver your baby is 14%. Without an epidural it is 7%.
- With an epidural, the second stage of labour (when your cervix is fully dilated) is longer and you are more likely to need medication (oxytocin) to make your contractions stronger.
- You have more chance of having low blood pressure.
- Your legs may feel weak while the epidural is working.
- You will find it difficult to urinate. You will probably need to have a tube passed into your bladder (a bladder catheter) to drain the urine.
- You may feel itchy.
- You may develop a slight fever.
- If you have higher doses of opioid through an epidural, your newborn baby may be more likely to need help with breathing (see reference 13) and you may have less chance of breastfeeding successfully (see reference 14).

Other risks

On average, having an epidural does not give you a higher risk of a headache. However, in around one in every 100 women who have an epidural the bag of fluid which surrounds their spinal cord is punctured by the epidural needle (this is called a 'dural puncture'). If this happens to you, you could get a severe headache that could last for days or weeks if it is not treated (see reference 15). If you do develop a severe headache, your anaesthetist should talk to you and give you advice about the treatment you could have. (For more details please see our '[Post dural puncture](#)' leaflet.)

The following information is based on the results of observational studies.

- The risks of epidurals and spinals are shown in the table below (see references 16 to 22).
- About one in every 24,000 women gets long-lasting nerve damage after an epidural, causing problems such as weakness in your leg or a feeling of tingling or numbness down one leg (reference 23). However, nerve damage after giving birth can happen whether you have an epidural or not (see reference 16) and is actually about five times more common without an epidural, with one in every 2,500 women being affected by it.
- There is no evidence to show that having an epidural while you are in labour causes the nerves in your spine to become permanently inflamed (that is, swollen and sore). See reference 22.

If you are worried about the risk of serious problems that might happen with an epidural, talk about this with your anaesthetist

Information App

This information is also available as an app on all types of smartphones. [You can download it from the OAA website](#)

Risks of having an epidural or spinal to reduce labour pain

Type of risk	How often does this happen?	How common is it?
Itching	One in every 3 to 10 women depending on the drug and dose used	Common
Significant drop in blood pressure	One in every 50 women	Occasional
Not working well enough to reduce labour pain so you need to use other ways of reducing the pain	One in every 10 women	Common
Not working well enough for a caesarean section so you need to have a general anaesthetic	One in every 20 women	Occasional
Severe headache	One in every 100 women (epidural)	Uncommon
Nerve damage (numb patch on a leg or foot, or having a weak leg)	One in every 500 women (spinal)	Quite rare
Effects lasting for more than 6 months	Temporary – one in every 1,000 to 2,000 women	Rare
Meningitis	Permanent – one in every 24,000 women	Very rare
	One in every 100,000 women	

References

- 1 Intrapartum care. Care of healthy women and their babies during childbirth. National Collaborating Centre for Women's and Children's Health. Commissioned by the National Institute for Health and Clinical Excellence. 2007 RCOG Press, London.
- 2 Waldenström U, Nilsson CA. Experience of childbirth in birth center care. A randomised controlled study. *Acta Obstetrica et Gynecologica Scandinavica* 1994; 73: 547-554.
- 3 Hodnett ED, Gates S, Hofmeyr GJ, Sakala C. Continuous support for women during childbirth. *Cochrane Database of Systematic Reviews* 2003, Issue 3. Article number: CD003766. Date of issue: 10.1002/14651858.CD003766.
- 4 Cluett ER, Nikodem VC, McCandlish RE, Burns EE. Immersion in water in pregnancy, labour and birth. *Cochrane Database of Systematic Reviews* 2002, Issue 2. Article number: CD000111. Date of issue: 10.1002/14651858.CD000111.pub2.
- 5 Smith CA, Collins CT, Cyna AM, Crowther CA. Complementary and alternative therapies for pain management in labour. *Cochrane Database of Systematic Reviews* 2006, Issue 4. Article number: CD003521. Date of issue: 10.1002/14651858.CD003521.pub2.
- 6 Wee MYK, Tuckley JP, Thomas P, Bernard S and Jackson D. The IDvIP trial: A two centre double blind randomised controlled trial comparing i.m. diamorphine and i.m. pethidine for labour. *International Journal of Obstetric Anaesthesia*. 2012;21(S1) S15.
- 7 Olofsson C, Ekblom A, Ekman-Ordeberg G, Hjelm A, Irestedt L. Lack of analgesic effect of systemically administered morphine or pethidine on labour pain. *British Journal of Obstetrics and Gynaecology* 1996;103:968-972.
- 8 Volmanen P, Akural E, Raudaskoski T, Ohtonen P, Alahuhta S. Comparison of remifentanyl and nitrous oxide in labour analgesia. *Acta Anaesthesiologica Scandinavica* 2005; 49: 453-458.
- 9 Volikas I, Butwick A. Maternal and neonatal side effects of remifentanyl PCA. *British Journal of Anaesthesia* 2005; 95: 504-509.
- 10 Anim-Somuah M, Smyth R, Howell C. Epidural versus non-epidural or no analgesia in labour. *Cochrane Database of Systematic Reviews* 2005, Issue 4. Article number: CD000331. Date of issue: 10.1002/14651858.CD000331.pub2.
- 11 Reynolds F, Sharma S, Seed PT. Analgesia in labour and funic acid-base balance: a meta-analysis comparing epidural with systemic opioid analgesia. *British Journal of Obstetrics and Gynaecology* 2002; 109: 1344-1353.
- 12 Russell R, Dundas R, Reynolds F. Long term backache after childbirth: prospective search for causative factors. *British Medical Journal* 1996; 312: 1384-1388.
- 13 COMET Study Group UK. Effect of low-dose mobile versus traditional epidural techniques on mode of delivery: a randomised controlled trial. *Lancet* 2001; 358: 19-23.
- 14 Beilin Y, Bodian CA, Weiser J, Hossain S, Arnold I, Feerman DE, Martin G, Holzman I. Effect of labor epidural analgesia with and without fentanyl on infant breast-feeding: a prospective, randomized, double-blind study. *Anesthesiology* 2005; 103: 1211-1217.
- 15 Sudlow C, Warlow C. Epidural blood patching for preventing and treating post-dural puncture headache. *Cochrane Database of Systematic Reviews* 2001, Issue 2. Article number: CD001791. Date of issue: 10.1002/14651858.CD001791.

- 16 Holdcroft A, Gibberd FB, Hargrove RL, Hawkins DF, Dellaportas CI. Neurological complications associated with pregnancy. *British Journal of Anaesthesia* 1995; 75: 522-526.
- 17 Jenkins K, Baker AB. Consent and anaesthetic risk. *Anaesthesia* 2003; 58: 962-984.
- 18 Jenkins JG, Khan MM. Anaesthesia for Caesarean section: a survey in a UK region from 1992 to 2002. *Anaesthesia* 2003; 58: 1114-1118.
- 19 Jenkins JG. Some immediate serious complications of obstetric epidural analgesia and anaesthesia: a prospective study of 145,550 epidurals. *International Journal of Obstetric Anesthesia* 2005; 14: 37-42.
- 20 Reynolds F. Infection a complication of neuraxial blockade. *International Journal of Obstetric Anesthesia* 2005; 14: 183-188.
- 21 Ruppen W, Derry S, McQuay H, Moore RA. Incidence of epidural hematoma, infection, and neurologic injury in obstetric patients with epidural analgesia/ anesthesia. *Anesthesiology* 2006; 105: 394-399.
- 22 Rice I, Wee MYK, Thomson K. Obstetric epidurals and chronic adhesive arachnoiditis. *British Journal of Anaesthesia* 2004; 92: 109-120.
23. Major complications of central neuraxial block: Report on the 3rd National Audit project of the Royal College of anaesthetist. 2009. RCOA.

LabourPains.com is the public website of the Obstetric Anaesthetists Association. This booklet was written by the Information for Mothers Subcommittee of the Obstetric Anaesthetists' Association. The subcommittee includes representatives from the National Childbirth Trust, the Royal College of Obstetricians and Gynaecologists, the Royal College of Midwives and patient representatives. This is based on the written leaflet, 4th edition, July 2013. © Obstetric Anaesthetists' Association 2013.